

On the Mellin Transform of the Coefficient Functions of $F_L(x, Q^2)$

Johannes Blümlein and Stefan Kurth

DESY-Zeuthen, Platanenallee 6, D-15735 Zeuthen, Germany

Abstract

The Mellin-transforms of the next-to-leading order Wilson coefficients of the longitudinal structure function are evaluated.

1 Introduction

For the study of the scaling violations of deep-inelastic scattering structure functions different techniques were developed [1, 2]. In most of the approaches the evolution equations of the parton densities are solved as integro-differential equations in x -space [1]. In the case of twist-2 operators the Mellin-transform $M[f](N)$, given by

$$M[f](N) = \int_0^1 dx x^{N-1} f(x) , \quad (1)$$

maps the convolutions between the parton densities, splitting functions, and coefficient functions into ordinary products, which leads to a considerable simplification of the problem [2]. Here N denotes the integer moment-index. The analytic continuation [3] of the functional $M[f](N)$ into the complex plane is later used to perform the inverse transformation to x -space. It appears useful to apply the Mellin-representation not only for the evolution equations of the parton densities but also to the structure functions themselves. Their representation in x -space is then obtained directly by a single inverse Mellin transform. For various next-to-leading order studies which were performed so far, as e.g. of the structure functions $F_2(x, Q^2)$, $g_1(x, Q^2)$, and $xF_3(x, Q^2)$, the coefficient functions were requested only in $O(\alpha_s)$ and the above procedure has been followed already. In the case of the structure function $F_L(x, Q^2)$ the NLO expressions [4–6] contain as well the Wilson coefficients in $O(\alpha_s^2)$. They were calculated in the x -representation in ref. [4] for the quarkonic and in ref. [5, 6] for the gluonic coefficient functions, where also the results for the quarkonic coefficient functions of ref. [4] were confirmed. A careful check of all results was performed in ref. [6] where a series of integer moments was evaluated both numerically and analytically and compared to the results of an independent calculation, ref. [7]. Previously the x -space expressions were used in various studies, see e.g. [8].

It is the aim of this note to shortly present the representation of the coefficient functions in Mellin- N space and their analytic continuation to allow also to evaluate the longitudinal structure function at next-to-leading order directly within the approach used in ref. [2].

2 The Wilson Coefficients in x space

The longitudinal structure function $F_L(x, Q^2)$ has the representation

$$F_L(x, Q^2) = x \left\{ C_{\text{NS}}(x, Q^2) \otimes f_{\text{NS}}(x, Q^2) + \delta_f \left[C_{\text{S}}(x, Q^2) \otimes \Sigma(x, Q^2) + C_{\text{g}}(x, Q^2) \otimes G(x, Q^2) \right] \right\} \quad (2)$$

in the case of pure photon exchange. The symbol $\otimes$ denotes the Mellin convolution

$$A(x, Q^2) \otimes B(x, Q^2) = \int_0^1 dx_1 \int_0^1 dx_2 \delta(x - x_1 x_2) A(x_1, Q^2) B(x_2, Q^2) . \quad (3)$$

The combinations of parton densities are

$$f_{\text{NS}}(x, Q^2) = \sum_{i=1}^{N_f} e_i^2 \left[q_i(x, Q^2) + \bar{q}_i(x, Q^2) \right] , \quad (4)$$

$$\Sigma(x, Q^2) = \sum_{i=1}^{N_f} \left[q_i(x, Q^2) + \bar{q}_i(x, Q^2) \right] . \quad (5)$$

$G(x, Q^2)$ denotes the gluon density, e_i the electric charge, and $\delta_f = (\sum_{i=1}^{N_f} e_i^2)/N_f$, with N_f the number of active flavors.

The coefficient functions $C_i(x, Q^2)$ are given by

$$\begin{aligned} C_{\text{NS}}(z, Q^2) &= a_s c_{L,q}^{(1)}(z) + a_s^2 c_{L,q}^{(2),\text{NS}}(z) \\ C_{\text{S}}(z, Q^2) &= a_s^2 c_{L,q}^{(2),\text{PS}}(z) \\ C_{\text{g}}(z, Q^2) &= a_s c_{L,g}^{(1)}(z) + a_s^2 c_{L,g}^{(2)}(z) , \end{aligned} \quad (6)$$

where $a_s = \alpha_s(Q^2)/(4\pi)$. For convenience we list as well the coefficient functions in x -space, since we will give the Mellin transforms of the individual contributing functions separately below. The leading order coefficient functions are given by [9]

$$c_{L,q}^{(1)}(z) = 4C_F z \quad (7)$$

$$c_{L,g}^{(1)}(z) = 8N_f z(1-z) . \quad (8)$$

In the $\overline{\text{MS}}$ scheme the NLO coefficient functions read [4–6]

$$\begin{aligned} c_{L,q}^{(2),\text{NS}}(z) &= 4C_F(C_A - 2C_F)z \left\{ 4 \frac{6 - 3z + 47z^2 - 9z^3}{15z^2} \ln z \right. \\ &\quad - 4\text{Li}_2(-z)[\ln z - 2\ln(1+z)] - 8\zeta(3) - 2\ln^2 z [\ln(1+z) + \ln(1-z)] \\ &\quad + 4\ln z \ln^2(1+z) - 4\ln z \text{Li}_2(z) + \frac{2}{5}(5 - 3z^2) \ln^2 z \\ &\quad - 4 \frac{2 + 10z^2 + 5z^3 - 3z^5}{5z^3} [\text{Li}_2(-z) + \ln z \ln(1+z)] \\ &\quad + 4\zeta(2) \left[\ln(1+z) + \ln(1-z) - \frac{5 - 3z^2}{5} \right] + 8\text{S}_{1,2}(-z) + 4\text{Li}_3(z) \\ &\quad + 4\text{Li}_3(-z) - \frac{23}{3} \ln(1-z) - \frac{144 + 294z - 1729z^2 + 216z^3}{90z^2} \Big\} \\ &\quad + 8C_F^2 z \left\{ \text{Li}_2(z) + \ln^2 z - 2\ln z \ln(1-z) + \ln^2(1-z) - 3\zeta(2) \right. \\ &\quad \left. - \frac{3 - 22z}{3z} \ln z + \frac{6 - 25z}{6z} \ln(1-z) - \frac{78 - 355z}{36z} \right\} \\ &\quad - \frac{8}{3} C_F N_f z \left\{ 2\ln z - \ln(1-z) - \frac{6 - 25z}{6z} \right\} , \end{aligned} \quad (9)$$

$$\begin{aligned} c_{L,q}^{(2),\text{PS}}(z) &= \frac{16}{9z} C_F N_f \left\{ 3(1 - 2z - 2z^2)(1-z) \ln(1-z) + 9z^2 [\text{Li}_2(z) + \ln^2 z - \zeta(2)] \right. \\ &\quad \left. + 9z(1 - z - 2z^2) \ln z - 9z^2(1-z) - (1-z)^3 \right\} , \end{aligned} \quad (10)$$

$$\begin{aligned} c_{L,g}^{(2)}(z) &= C_F N_f \left\{ 16z [\text{Li}_2(1-z) + \ln z \ln(1-z)] \right. \\ &\quad + \left(-\frac{32}{3}z + \frac{64}{5}z^3 + \frac{32}{15z^2} \right) [\text{Li}_2(-z) + \ln z \ln(1+z)] + (8 + 24z - 32z^2) \ln(1-z) \\ &\quad - \left(\frac{32}{3}z + \frac{32}{5}z^3 \right) \ln^2 z + \frac{1}{15} \left(-104 - 624z + 288z^2 - \frac{32}{z} \right) \ln z \\ &\quad + \left(-\frac{32}{3}z + \frac{64}{5}z^3 \right) \zeta(2) - \frac{128}{15} - \frac{304}{5}z + \frac{336}{5}z^2 + \frac{32}{15z} \Big\} \\ &\quad + C_A N_f \left\{ -64\text{Li}_2(1-z) + (32z + 32z^2) [\text{Li}_2(-z) + \ln z \ln(1+z)] \right. \end{aligned}$$

$$\begin{aligned}
& +(16z - 16z^2) \ln^2(1 - z) + (-96z + 32z^2) \ln z \ln(1 - z) \\
& + \left(-16 - 144z + \frac{464}{3}z^2 + \frac{16}{3z} \right) \ln(1 - z) + 48z \ln^2 z + (16 + 128z - 208z^2) \ln z \\
& + 32z^2 \zeta(2) + \frac{16}{3} + \frac{272}{3}z - \frac{848}{9}z^2 - \frac{16}{9z} \Big\} ,
\end{aligned} \tag{11}$$

with $C_A = N_c = 3, C_F = (N_c^2 - 1)/(2N_c) = 4/3$. The corresponding expressions in the DIS scheme are given in [6]. The class of basic functions is the same for both schemes.

3 The Mellin Transform

For most of the functions $f_i(x)$ contributing to eqs. (7–11) the Mellin transforms may be evaluated straightforwardly¹. They are listed in Table 1 for the individual functions. In these cases infinite (or finite) sums occurring may be traced back to representations being based on the functions

$$\psi^{(k)}(z) = \frac{1}{\Gamma(z)} \frac{d^k}{dz^k} \Gamma(z) , \tag{12}$$

which represent the sums

$$S_n(N) = \sum_{k=1}^N \frac{1}{k^n} , \tag{13}$$

with

$$S_1(N) = \psi(N+1) + \gamma_E , \tag{14}$$

and

$$S_n(N) = \frac{(-1)^{n-1} \psi^{(n-1)}(N+1)}{\Gamma(n)} + \zeta(n), \quad n > 1 . \tag{15}$$

Here $\gamma_E = -\psi(1)$ is the Mascheroni constant and $\zeta(n) = \sum_{k=1}^{\infty} (1/k^n)$ denotes the Riemann ζ -function. One further may use Euler's relation

$$\text{Li}_2(1 - z) = -\text{Li}_2(z) - \ln z \ln(1 - z) + \zeta(2) \tag{16}$$

to reduce the number of functions in the above expressions. Similarly it is convenient to combine the functions $\text{Li}_2(-x) + \ln x \ln(1 + x)$ whenever they occur with the same coefficient to

$$\text{Li}_2(-x) + \ln x \ln(1 + x) = -\frac{1}{2} \tilde{\Phi}(x) + \frac{1}{4} \ln^2 x - \frac{\zeta(2)}{2} , \tag{17}$$

since the Mellin transform of the function

$$\tilde{\Phi}(x) = \int_{x/(1+x)}^{1/(1+x)} \frac{dz}{z} \ln \left(\frac{1-z}{z} \right) \tag{18}$$

turns out to be simpler than that of either $\text{Li}_2(-x)$ or $\ln x \ln(1 + x)$. All these functions can be traced back to expressions which are built solely out of the functions $\psi^{(k)}(z)$, see Table 1.

¹Useful relations can be found in refs. [10, 11].

No.	$f(z, r)$	$M[f](N)$
1	z^r	$\frac{1}{N+r}$
2	$z^r \ln^n z$	$\frac{(-1)^n}{(N+r)^{n+1}} \Gamma(n+1)$
3	$z^r \ln(1-z)$	$-\frac{S_1(N+r)}{N+r}$
4	$z^r \ln(1+z)$	$\frac{(-1)^{N+r-1}}{N+r} \left[-S_1(N+r) + \frac{1+(-1)^{N+r-1}}{2} S_1\left(\frac{N+r-1}{2}\right) \right. \\ \left. + \frac{1-(-1)^{N+r-1}}{2} S_1\left(\frac{N+r}{2}\right) \right] + [1+(-1)^{N+r+1}] \frac{\ln(2)}{N+r}$
5	$z^r \ln^2(1-z)$	$\frac{S_1^2(N+r) + S_2(N+r)}{N+r}$
6	$z^r \ln z \ln(1-z)$	$\frac{S_1(N+r)}{(N+r)^2} + \frac{1}{N+r} [S_2(N+r) - \zeta(2)]$
7	$z^r \ln^2 z \ln(1-z)$	$\frac{2}{N+r} \left[\zeta(3) + \frac{\zeta(2)}{N+r} - \frac{S_1(N+r)}{(N+r)^2} - \frac{S_2(N+r)}{N+r} - S_3(N+r) \right]$
8	$z^r \ln^2 z \ln(1+z)$	$2 \frac{(-1)^{N+r}}{N+r} \left[\frac{S_1(N+r)}{(N+r)^2} + \frac{S_2(N+r)}{N+r} + S_3(N+r) - \frac{\zeta(2)}{2(N+r)} - \frac{3\zeta(3)}{4} \right] \\ + \frac{1+(-1)^{N+r-1}}{2(N+r)} \left[\frac{2}{(N+r)^2} S_1\left(\frac{N+r-1}{2}\right) + \frac{1}{N+r} S_2\left(\frac{N+r-1}{2}\right) \right. \\ \left. + \frac{1}{2} S_3\left(\frac{N+r-1}{2}\right) + \frac{4 \ln 2}{(N+r)^2} \right] - \frac{1-(-1)^{N+r-1}}{2(N+r)} \left[\frac{2}{(N+r)^2} S_1\left(\frac{N+r}{2}\right) \right. \\ \left. + \frac{1}{N+r} S_2\left(\frac{N+r}{2}\right) + \frac{1}{2} S_3\left(\frac{N+r}{2}\right) \right]$
9	$z^r \text{Li}_2(z)$	$\frac{1}{N+r} \left[\zeta(2) - \frac{S_1(N+r)}{N+r} \right]$
10	$z^r \text{Li}_2(z) \ln z$	$\frac{1}{(N+r)^2} \left[-2\zeta(2) + \frac{2S_1(N+r)}{N+r} + S_2(N+r) \right]$

Table 1 : Mellin transforms of basic functions contributing to the Wilson coefficients of $F_L(x, Q^2)$ in NLO.

No.	$f(z, r)$	$M[f](N)$
11	$z^r \tilde{\Phi}(z)$	$\frac{1}{(N+r)^3} + 2 \frac{(-1)^{N+r}}{N+r} \left[S_2(N+r) - \zeta(2) \right]$ $- \frac{1 + (-1)^{N+r}}{2(N+r)} \left[S_2 \left(\frac{N+r}{2} \right) - \zeta(2) \right]$ $+ \frac{1 - (-1)^{N+r}}{2(N+r)} \left[S_2 \left(\frac{N+r-1}{2} \right) - \zeta(2) \right]$
12	$z^r \text{Li}_2(-z) \ln z$	$\frac{(-1)^{N+r-1}}{(N+r)^2} \left[\frac{2S_1(N+r)}{N+r} + S_2(N+r) \right]$ $- \frac{1 + (-1)^{N+r-1}}{2(N+r)^2} \left[\frac{2}{N+r} S_1 \left(\frac{N+r-1}{2} \right) + \frac{1}{2} S_2 \left(\frac{N+r-1}{2} \right) + \frac{4 \ln 2}{N+r} \right]$ $+ \frac{1 - (-1)^{N+r-1}}{2(N+r)^2} \left[\frac{2}{N+r} S_1 \left(\frac{N+r}{2} \right) + \frac{1}{2} S_2 \left(\frac{N+r}{2} \right) + \zeta(2) \right]$
13	$z^r \text{Li}_3(z)$	$\frac{1}{N+r} \left[\zeta(3) - \frac{\zeta(2)}{N+r} + \frac{S_1(N+r)}{(N+r)^2} \right]$
14	$z^r \text{Li}_3(-z)$	$(-1)^{N+r-1} \frac{S_1(N+r)}{(N+r)^3} - \frac{1 + (-1)^{N+r-1}}{2(N+r)^3} \left[S_1 \left(\frac{N+r-1}{2} \right) + 2 \ln 2 \right]$ $+ \frac{1 - (-1)^{N+r-1}}{2(N+r)^3} S_1 \left(\frac{N+r}{2} \right) + \frac{\zeta(2)}{2(N+r)^2} - \frac{3\zeta(3)}{4(N+r)}$

Table 1 continued

To the non-singlet coefficient function also the functions

$$\ln(x) \ln^2(1+x), \quad S_{1,2}(-x), \quad \text{and} \quad \text{Li}_2(-x) \ln(1+x) \quad (19)$$

contribute, which lead to different sums. The Mellin transform of $\ln(x) \ln^2(1+x)$ can be obtained by a serial expansion

$$\begin{aligned}
\int_0^1 dx x^{N-1} \ln x \ln^2(1+x) &= \frac{2\gamma_E}{N^2} \left[\ln 2 - \frac{1}{2} \psi \left(\frac{N}{2} + 1 \right) + \frac{1}{2} \psi \left(\frac{N+1}{2} \right) \right] \\
&+ \frac{\gamma_E}{2N} \left[\psi' \left(\frac{N}{2} + 1 \right) - \psi' \left(\frac{N+1}{2} \right) \right] - 2 \sum_{k=1}^{\infty} \frac{(-1)^k \psi(k)}{k(N+k)^2} . \quad (20)
\end{aligned}$$

The infinite sum in eq. (20) is easily evaluated by a recursive algorithm. For numerical applications one may use as well representations of $\log(1+x)$ in the range $x \in [0, 1]$, cf. [12], as, e.g.,

$$\ln(1+x) \simeq \sum_{k=1}^8 a_k x^k, \quad (21)$$

which are as accurate as 10^{-8} . The coefficients a_k read

$$\begin{aligned}
a_1 &= 0.9999964239 & a_2 &= -0.4998741238 & a_3 &= 0.3317990258 & a_4 &= -0.2407338084 \\
a_5 &= 0.1676540711 & a_6 &= -0.0953293897 & a_7 &= 0.0360884937 & a_8 &= -0.0064535442 .
\end{aligned}$$

The Mellin transform of $\ln x \ln^2(1+x)$ is then given by

$$\int_0^1 dx x^{N-1} \ln x \ln^2(1+x) \simeq - \sum_{k=2}^{16} \frac{b_k}{(N+k)^2} , \quad (22)$$

where the coefficients b_k are obtained by taking the square of the polynomial in eq. (21).

We furthermore observe that $S_{1,2}(-x)$ and $\text{Li}_2(-x) \ln(1+x)$ may be combined using the integral representation, cf. [11]²,

$$F_1(x) = \frac{1}{x} \left\{ \ln(1+x) \text{Li}_2(-x) - \zeta(2) + S_{1,2}(-x) - 2\text{Li}_3(-x) \right\} = \int_0^1 dy \frac{\ln(y) \ln(1-y)}{1+xy} . \quad (23)$$

Since the functions $S_{1,2}(-x)$ and $\text{Li}_2(-x) \ln(1+x)$ occur with the same weight factor they can be dealt with together with the help of the representation eq. (23). In evaluating the Mellin-transform the x -integral yields

$$\sum_{k=0}^{\infty} (-1)^k \frac{y^k}{N+k} \equiv \Phi(-y, 1, N). \quad (24)$$

Here the function $\Phi(y, a, b)$, ref. [13], is related to the generalized Riemann ζ -function $\zeta(c, n)$ [14]. Calculating the y -integral by using eq. (6) of table 1, one finally obtains

$$M[F_1](N) = \sum_{k=0}^{\infty} \frac{(-1)^k}{N+k} \left[\frac{\psi(2+k) + \gamma_E}{(k+1)^2} - \frac{\psi'(2+k)}{k+1} \right]. \quad (25)$$

The representation eq. (25) is fastly converging since for large values of k , $\psi(k) \sim \ln(k)$ and $\psi'(k) \sim 1/k$. As in the case of the Mellin transforms of other functions emerging in the x -space representation of the different coefficient and splitting functions the poles in eqs. (20) and (25) are situated at the non-positive integers.

We finally would like to comment on the analytic continuation of the sums

$$\tilde{S}_n(N) = \sum_{k=1}^N \frac{(-1)^k}{k^n} S_1(k) , \quad n \geq 2. \quad (26)$$

which emerge in some of the Mellin-transforms. $\tilde{S}_2(N)$ contributes, e.g., to the NLO anomalous dimensions. By using the relation

$$c_{k,n} = \sum_{l=1}^k \frac{(-1)^l}{l^n} = - \left\{ \left(1 - \frac{1}{2^{n-1}} \right) \zeta(n) + \frac{(-1)^{k+n-1}}{2^n \Gamma(n)} \left[\psi^{(n-1)} \left(\frac{k+1}{2} \right) - \psi^{(n-1)} \left(\frac{k}{2} + 1 \right) \right] \right\} \quad (27)$$

one may express $\tilde{S}_k(N)$ as

$$\begin{aligned} \tilde{S}_n(N) &= (-1)^N \frac{S_1(N)}{N^n} - \sum_{l=2}^n (-1)^l \zeta(l) \left\{ \left(1 - \frac{1}{2^{n-l}} \right) \zeta(n-l+1) \right. \\ &\quad \left. - (-1)^N \frac{(-1)^{n-l}}{2^{n-l+1} \Gamma(n-l+1)} \left[\psi^{(n-l)} \left(\frac{N}{2} \right) - \psi^{(n-l)} \left(\frac{N+1}{2} \right) \right] \right\} \\ &\quad - (-1)^n \left[\zeta(n) \ln 2 - (-1)^N \int_0^1 dx \frac{x^{N-1} \text{Li}_n(x)}{1+x} - \int_0^1 dx \frac{\text{Li}_n(x)}{1+x} \right], \end{aligned} \quad (28)$$

²Note a misprint in eq. (3.12.22) of ref. [11]. The factor in front of $\text{Li}_3(-b/a)$ should be 2.

where the Mellin-transform of $\text{Li}_n(x)/(1+x)$ is given by

$$M[F_2](N) = \int_0^1 dx \frac{\text{Li}_n(x)}{1+x} x^{N-1} = \sum_{k=1}^{\infty} \frac{(-1)^k}{N+k} c_{k,n} . \quad (29)$$

The series eq. (29) converges rather fast for $n > 2$. For the special case $n = 2$, being dealt with before in ref. [15], one obtains from (28)

$$\tilde{S}_2(N) = -\frac{5}{8}\zeta(3) + (-1)^N \left[\frac{S_1(N)}{N^2} - \frac{\zeta(2)}{2} G(N) + \int_0^1 dx x^{N-1} \frac{\text{Li}_2(x)}{1+x} \right], \quad (30)$$

with

$$\int_0^1 dx \frac{\text{Li}_2(x)}{1+x} = \zeta(2) \ln 2 - \frac{5}{8}\zeta(3) \quad \text{and} \quad G(N) = \psi\left(\frac{N+1}{2}\right) - \psi\left(\frac{N}{2}\right) . \quad (31)$$

For $n = 2$ the series eq. (29) converges slowly since the modulus of the expansion coefficients $c_{k,2}$ approach $\zeta(2)/2 = \pi^2/12$ as $k \rightarrow \infty$ and the series is essentially logarithmic. One may, however, rewrite $M[F_2](N)$ as

$$M[F_2](N) = \zeta(2) \ln(2) - \int_0^1 dx x^{N-2} \ln(1+x) [(N-1)\text{Li}_2(x) - \ln(1-x)] . \quad (32)$$

Again the function $\ln(1+x)$ can be represented by eq. (21) yielding

$$M[F_2](N) = \zeta(2) \ln(2) - \sum_{k=1}^8 a_k \left\{ (N-1) \left[\frac{\zeta(2)}{N+k-1} - \frac{\psi(N+k) + \gamma_E}{(N+k-1)^2} \right] + \frac{\psi(N+k) + \gamma_E}{N+k-1} \right\} . \quad (33)$$

which holds at an accuracy of better than $4 \cdot 10^{-7}$ for $N \in [1, 20]$. Another approximate expression for $M[F_2](N)$ was given in [15].

Acknowledgement. We would like to thank W. van Neerven, J. Vermaseren, F. Ynduráin, and A. Vogt for useful discussions.

References

- [1] For older approaches see refs. [46,48] in : J. Blümlein, *Surv. High Energy Phys.* **7** (1994) 181;
M. Virchaux and A. Ouraou, *DphPE* 87–15;
M. Virchaux, *Thèse, Université Paris–7*, 1988;
A. Ouraou, *Thèse, Université Paris–11*, 1988;
M. Botje, *QCDNUM15: A fast QCD evolution Program*, to appear;
C. Pascaud and F. Zomer, *H1 Note: H1–11/94–404*;
H. Lai et al., *CTEQ collaboration*, *Phys. Rev.* **D66** (1997) 1280, and references therein;
A.D. Martin, R. Roberts, M. Ryskin, and W. Stirling, *hep-ph/9612449* and references therein.
- [2] M. Diemoz, F. Ferroni, E. Longo, and G. Martinelli, *Z. Phys.* **C39** (1988) 21;
M. Glück, E. Reya, and A. Vogt, *Z. Phys.* **C48** (1990) 471.
- [3] For a uniqueness theorem see, E. Carlson, *Thesis*, Univ. Uppsala, 1914;
E.C. Titchmarsh, *Theory of Functions*, (Oxford University Press, Oxford, 1939), Chapt. 9.5.

- [4] J. Sanchez Guillén et al., Nucl. Phys. **B353** (1991) 337.
- [5] E. Zijlstra and W. van Neerven, Phys. Lett. **B273** (1991) 476.
- [6] E. Zijlstra and W. van Neerven, Nucl. Phys. **B383** (1992) 525;
E. Zijlstra, PhD Thesis, Leiden University, 1993.
- [7] S. Larin and J. Vermaseren, Z. Phys. **C57** (1993) 93.
- [8] J. Blümlein, Nucl. Phys. **B** (Proc. Suppl.) **B,C 39** (1994) 22, hep-ph/9408337;
J. Blümlein and S. Riemersma, Proc. of the 1996 Hera Physics Workshop, eds. A. DeRoeck, G. Ingelman, and R. Klanner, (DESY, Hamburg, 1996), **Vol. 1**, p. 82, hep-ph/9609364;
H1 Collaboration, C. Adloff et al., Phys. Lett. **B393** (1997) 452;
J. Blümlein and A. Vogt, hep-ph/9707488, DESY 97-143.
- [9] A. Zee, F. Wilczek, and S.B. Treiman, Phys. Rev. **D10** (1974) 2881.
- [10] A. González-Arroyo, C. López, and F.J. Ynduráin, Nucl. Phys. **B153** (1979) 161;
F.J. Ynduráin, *The Theory of Quark and Gluon Interactions*, (Springer, Berlin, 1993).
- [11] A. Devoto and D.W. Duke, La Riv. Nuovo Cim. **7** (1984) 1.
- [12] C. Hastings jr., *Approximations for Digital Computers*, (Princeton University Press, Princeton NJ, 1955), p. 180;
M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions*, (NBS, Washington, 1964), 4.1.44.
- [13] A. P. Prudnikov, Yu.A. Bryuchkov, and O.I. Marichev, *Integraly i Ryady: Elementarnye Funkzii*, (Nauka, Moscow, 1984) (in Russian); English Edition: (Gordon and Breach, Birmingham, 1986).
- [14] A. P. Prudnikov, Yu.A. Bryuchkov, and O.I. Marichev, *Integraly i Ryady: Dopolnitel'nye Glawy*, (Nauka, Moscow, 1986) (in Russian); English Edition: (Gordon and Breach, Birmingham, 1990).
- [15] M. Glück, E. Reya, and A. Vogt, Phys. Rev. **D45** (1992) 3986.